

bas condiciones permanece aún desconocida.

## BIBLIOGRAFIA

1. Kuo, T., Shih, L.Y., Chan, H.L.: Kimura's disease, involvement of regional lymph nodes and distinction from angiolympoid hyperplasia with eosinophilia. Am. J. Surg. Pathol. 1988; 12: 843-54.
2. Okudaira, H., Hongo, O., Ogita, T., et al.: Serum Ig E antibody levels in patients with bronchial asthma, atopic, eosinophilic granulomas diseases. Ann. Allergy. 1983; 50: 51-4.
3. Hui, P.K., Chan, J.K., Kung, L.T., Gwi, E.: Lymphadenopathy of Kimura disease. A. J. Surg. Pathol. 1989; 13: 177-18.
4. Kum, I.T., Gibson, J.B., Bannatyne, P.M.: Kimura's disease: a clinicopathological study of 21 cases and its distinction from angiolympoid hyperplasia with eosinophilia. Pathology. 1984; 16: 39-44.
5. Chan, J.C.K., Hui, P.K., et al.: Epithelioid hemangioma (angiolympoid hyperplasia with eosinophilia) and Kimura's disease in Chinese. Histopathology. 1989; 15: 557-74.
6. Kung, I.T.M., Chan, J.K.C.: Kimura's disease or Kim's disease? (Letter to the editor). Am. J. Surg. Pathol. 1988; 12: 804-5.
7. Dannaker, C., Piacquadio, D., Willoughby, C.B., et al.: Histiocytoid hemangioma: a disease spectrum. J. Am. Acad. Dermatol. 1989; 21: 404-9.
8. Urabe, A., Tsuneyoshi, M., Enjoji, M.: Epithelioid hemangioma versus Kimura's disease: a comparative clinicopathologic study. Am. J. Surg. Pathol. 1987; 11: 758-66.
9. Chun, S.I., Jih, G.: Kimura's disease and angiolympoid hyperplasia with eosinophilia: clinical and histopathologic differences. J. Am. Acad. Dermatol. 1992; 27: 954-8.
10. Welis, G.C., Whimster, I.W.: Subcutaneous angiolympoid hyperplasia with eosinophilia. Br. J. Dermatol. 1982; 118: 166-70.
11. Rosal, J., Angiolympoid hyperplasia with eosinophilia of the skin: its nosological position in the spectrum of histiocytoid hemangioma. Am. J. Dermatopathol. 1982; 4: 175-84.
12. Googe, P.B., Harris, N.L., Mihm, M.C.: Kimura's disease and angiolympoid hyperplasia with eosinophilia: two distinct histopathological entities. Am. J. Pathol. 1987; 14: 263-71.
13. Henry, P., Burnett, J.: Angiolympoid hyperplasia with eosinophilia. Arch. Dermatol. 1987; 114: 1168-72.
14. Fetsch, J.F., Weiss, S.W.: Observations concerning the pathogenesis of epithelioid hemangioma (angiolympoid hyperplasia). Moderm. Pathol. 1991; 40: 44955.

Anderson C.R. et al. Well's  
**Syndrome in childhood: Case report and review of the literature.**  
JAAD. Nov. 1995; 33: 857-64.

## CELULITIS EOSINOFILICA

Se trata de una dermatosis inflamatoria recurrente de patogénesis desconocida. Las lesiones cutáneas inicialmente semejan celulitis bacteriana e histopatológicamente presentan eosinofilia dérmica, histiocitos y figuras en llama. Ha sido descrito principalmente en adultos y sólo 17 casos en niños, con un rango de edad entre 3 sem. a 14 años, la mayoría varones, pueden haber lesiones ampollares, tiene curso benigno, con eosinofilia periférica, siendo el único tratamiento efectivo los esteroides sistémicos a altas dosis.

Dra. María Paola Leonardis L